

LOG OUT | SEARCH | MY PRODUCTS | CONTACT US | USER GUIDE

27th Edition AC ISC GRG MCM RFC HC CCG TC BHG PIP MCR Date of Service mm/dd/y

Quick Search irf Search

Recovery Facility Care > Optimal Recovery Guidelines > Subacute/Skilled Nursing Facility (SNF) > General Surgery > Bowel Surgery: Small Intestine Resection (S-5250)

Bowel Surgery: Small Intestine Resection

MCG Health Recovery Facility Care

27th Edition

ORG: S-5250 (RFC) Link to Codes

Print View

- · Clinical Indications for Admission to Recovery Facility
- · Alternative Levels of Care
- Length of Stay
- · Evaluation and Treatment
 - · General Treatment Course
 - Extended Stay
 - · Patient Education
- Discharge Planning
- Quality Measures
- References
- Footnotes
- Codes

Clinical Indications for Admission to Recovery Facility

Return to top of Bowel Surgery: Small Intestine Resection - RFC

- Skilled nursing facility (SNF) or subacute facility care is/was needed for appropriate care of patient because of ALL of the following:
 - There are no acute hospital care needs.(1)
 - The patient has Intense and complex care needs that make recovery facility care safer and more practical than attempting care at a lower level and **ALL** of the following[A][B](4):
 - These care needs include the multiple components of care that are delivered by skilled professionals at a recovery facility.[C][D][E]
 - There is a plan to provide **ALL** of the following(14):
 - Care plan management and evaluation to meet patient needs, achieve treatment goals, and ensure medical safety(15)
 - Observation and assessment of patient's changing condition to evaluate need for treatment modification or for additional procedures until condition stabilized
 - Education services to teach patient self-maintenance or to teach caregiver patient care(16)(17)
 - Skilled care treatments daily (or more frequent), including 1 or more of the following:
 - Nursing treatments needed for **1 or more** of the following(18):
 - Intravenous (IV) infusion, IV injection, or intramuscular (IM) injection[F](1)(16)(19)
 - Tube feeding (ie, gastrostomy tubes, jejunostomy tubes, percutaneous endoscopic gastrostomy (PEG) tubes, nasogastric tubes) required because patient needs feeding to supply at least 26% of daily calories and at least 501 mL of daily fluids(16)(20)
 - Wound management that requires dressing changes with prescription medication or clean technique and treatment for 1 or more of the following(1):
 - · Surgical wound complications
 - Patient care training and assistance for **1 or more** of the following(21):
 - Exercise program (eg, range of motion, pulmonary, cardiac)(22)